

LC-MS/MS Profiling-Based Secondary Metabolite Screening of *Myxococcus xanthus*

Kim, Jiyoung¹, Jung Nam Choi¹, Pil Kim², Dai-Eun Sok³, Soo-Wan Nam⁴, and Choong Hwan Lee^{1*}

¹Functional Metabolomics Laboratory, Division of Bioscience and Biotechnology, Konkuk University, Seoul 143-701, Korea

²Department of Biotechnology, Catholic University of Korea, Bucheon 420-743, Korea

³Chungnam National University, Daejeon 305-764, Korea

⁴Department of Biotechnology and Bioengineering/Department of Biomaterial Control, Dong-Eui University, Busan 614-714, Korea

Received: November 19, 2007 / Accepted: July 31, 2008

Myxobacteria, Gram-negative soil bacteria, are a well-known producer of bioactive secondary metabolites. Therefore, this study presents a methodological approach for the high-throughput screening of secondary metabolites from 4 wild-type *Myxococcus xanthus* strains. First, electrospray ionization mass spectrometry (ESI-MS) was performed using extracellular crude extracts. As a result, 22 metabolite peaks were detected, and the metabolite profiling was then conducted using the m/z value, retention time, and MS/MS fragmentation pattern analyses. Among the peaks, one unknown compound peak was identified as analogous to the myxalamid A, B, and C series. An analysis of the tandem mass spectrometric fragmentation patterns and HR-MS identified myxalamid K as a new compound derived from *M. xanthus*. In conclusion, LC-MS/MS-based chemical screening of diverse secondary metabolites would appear to be an effective approach for discovering unknown microbial secondary metabolites.

Keywords: Metabolite profiling, LC-MS/MS, *Myxococcus xanthus*, myxalamid

Microorganisms are an abundant source of new useful natural products [10], and myxobacteria have been shown to be prolific producers of a variety of bioactive secondary metabolites, including antibacterial and antifungal compounds [6, 13].

Recently, the metabolomics approach has been used to classify metabolites based on metabolite profiling studies, allowing rapid analyses of complex data and the determination of novel compounds [15, 17]. The increased interest in metabolite profiling has also arisen from the potential for more comprehensive metabolite analyses using liquid

chromatography mass spectrometry (LC-MS) technology [1]. The most important advantages of LC-MS are its high sensitivity and high-throughput in combination with the possibility to confirm the identity of the components present in complex biological samples, along with the detection and identification of unknown and unexpected compounds [7, 16].

Accordingly, to screen diverse secondary metabolites, wild-type *Myxococcus* strains were collected from the MicroBank (<http://www.microbank.re.kr>) of the Korea Research Institute of Bioscience and Biotechnology (KRIBB) [11]. The 4 selected strains were cultivated on CY agar plates for 3–4 days at 30°C [14]. The basal medium for the metabolite profiling was cultivated in 100 ml of CY media for 7 days at 30°C on a rotary shaker at 150 rpm. Amberlite XAD-16 resin (2%, v/v) was added to promote the productivity of the active substance after 3 days of cultivation [2, 12]. At the end of the cultivation period, the XAD-16 adsorber resin was separated from the culture broth, followed by washing with water for 20 min at 30°C and elution twice with 20 ml of methanol. The methanol extracts were then evaporated under the simultaneous addition of water to a final volume of 50 ml. Thereafter, the water phase was extracted with 10 ml of ethyl acetate, and the organic phase concentrated. Finally, the residues were dissolved with 80% methanol (3 mg/ml) and analyzed by LC-MS. In this study, the LC-MS was performed using a Finnigan LCQ Advantage MAX ion trap mass spectrometer (Thermo Electron Co., U.S.A.) equipped with an electrospray ionization (ESI) source. The HPLC separation was performed using a Finnigan Surveyor Modular HPLC System (Thermo Electron Co., U.S.A.) with a YMC-Pack Pro C18 column (2.1×50 mm, 5 μm), where the system was operated using Xcalibur software (version 1.3 SP2, Thermo Electron). The mobile phases A and B were water and acetonitrile with 0.1% formic acid, respectively, and the gradient elution at a flow rate of 0.2 ml/min was performed as

*Corresponding author

Phone: 82-2049-6177; Fax: 82-2-455-4291;

E-mail: chlee123@konkuk.ac.kr

follows: 0–30 min for 5–100% B (linear gradient), 30–40 min for 100% B (isocratic), followed by 40–50 min of 5% B (isocratic). The full-scan mass spectra were obtained within a range of m/z 50–1,500 using three microscans. Utilizing the capacity of the LCQ mass spectrometer to provide continuous polarity switching, data acquisition was conducted in positive and negative modes. The data-dependent tandem mass spectrometry (MS/MS) experiments were controlled using the menu-driven software provided with the Xcalibur system. All the experiments were performed under automatic gain control conditions. The high-resolution mass spectrometer (HR-MS) was measured in methanol with a PEG400 on a JMS-700 Mstation (JEOL Ltd.) consisting of a high-resolution two-sector mass spectrometer using the FAB ionization method from the Korea Basic Science Institute (KBSI). The ^1H NMR 500 MHz was measured in CD_3OD on a Varian UI500 with equipment analysis support from KBSI.

The LC-MS analytical data were optimized using a background subtraction technique of chromatography with the metabolite ID 2.0 software system. The principle of this method is to reduce the background, such as noise and fault peaks, to subtract the medium components from the sample extracts (Fig. 1). In the subsequent optimization for the chromatography and spectrum analyses, data containing more real secondary metabolites were observed, along with more ions present in the processed mass spectra. After exporting the process, the LC-MS data were manually




Fig. 1. Background and medium subtracted positive ion chromatogram of *M. xanthus* KYC1427.

A. Chromatogram of medium; B. Culture extracts of KYC1427; C. Subtracted (B)-(A) using Metabolite ID 2.0.

sorted to list such information as the retention time, m/z values for $[\text{M}+\text{H}]^+$, MS/MS fragmentation pattern, and UV spectra from base peak chromatograms (Table 1). Each compound was then identified from reference secondary metabolites already known to be produced by *M. xanthus*.

Table 1. Screening of secondary metabolites found in crude extracts of *Mycococcus* strains.

No.	Metabolites	Rt (min)	$[\text{M}+\text{H}]^+$ (m/z)	Fragmented ion in MS2 spectra (m/z)	ATCC25232	KYC1288	KYC1427	KYC1461	Ref.
1	Myxoachelin B	7.09	404	387,251	*	*	*	*	[8]
2	Cittilin A	8.63	631	614,603,586,558,518,490,462,445				*	[13]
3	Myxoachelin A	10.59	405	387,269	*	*			[8]
4	Unknown	10.68	446	428,386,310			*	*	
5	Unknown	11.27	460	442,386,324			*		
6	Unknown	11.79	419	401,283	*				
7	DKxanthene-534	12.08	535	518,500,444,416,399,268,251		*			[9]
8	DKxanthene-518	12.85	519	502,400,383,320,268,251	*	*	*	*	[9]
9	Unknown	13.73	545	528,426,408,294		*	*	*	
10	Unknown	14.01	552	534,416	*				
11	Unknown	14.85	536	518,503,426,400,382			*		
12	Unknown	15.23	488	470,401,352	*				
13	Unknown	16.33	404	387,267,281			*		
14	Unknown	16.35	502	484,366	*				
15	Unknown	17.19	516	498,472,442,366	*		*		
16	Unknown	19.77	578	503,489,152,419,416	*				
17	Myxalamid C	20.34	388	370,313,295,290			*		[4]
18	Myxovirescin Bor N1	21.21	622	594,341,286		*			[3]
19	Myxovirescin A2 or T1	21.81	624	579,473		8			[3]
20	Myxalamid B	21.58	402	384,327,309,290,267,215	*		*		[4]
21	Myxalamid A	22.32	416	398,341,323,290,267,215	*	*	*	*	[4]
22	Myxalamid K (New)	23.72	430	412,355,337,304,267,215	*	*	*	*	

files. As shown in the present study, the LC-MS/MS-based metabolite profiling of *M. xanthus* secondary metabolites would appear to be a useful technique for distinguishing between known and unknown compounds, and for screening novel compounds without large amounts of cultures. Furthermore, the method saves time and minimizes the repeated discovery of known metabolites.

Acknowledgments

This work was supported by the Korea Science and Engineering Foundation (No. R01-2007-000-20231-0) and Microbial Genomics and Application Center funded by the Korea government (MEST).

REFERENCES

- Bino, R. J., R. D. Hall, O. Fiehn, J. Kopka, K. Saito, J. Draper, *et al.* 2004. Potential of metabolomics as a functional genomics tool. *Trends Plant Sci.* **9**: 418–424.
- Frykman, S., H. Tsuruta, J. Galazzo, and P. Licari. 2006. Characterization of product capture resin during microbial cultivations. *J. Ind. Microbiol. Biotechnol.* **33**: 445–453.
- Gerth, K., H. Irschik, H. Reichenbach, and W. Trowitzsch. 1982. The myxovirescins, a family of antibiotics from *Myxococcus virescens* (Myxobacterales). *J. Antibiot.* **35**: 1454–1459.
- Gerth, K., R. Jansen, G. Reifenhahl, G. Höfle, H. Irschik, B. Kunze, H. Reichenbach, and G. Thierbach. 1983. The myxalamids, new antibiotics from *Myxococcus xanthus* (Myxobacterales). I. Production, physico-chemical and biological properties and mechanism of action. *J. Antibiot.* **36**: 1150–1156.
- Jansen, R., G. Reifenhahl, K. Gerth, H. Reichenbach, and G. Höfle. 1983. Myxalamide A, B, C, and D, eine gruppe homologer antibiotika aus *Myxococcus xanthus* Mxx12 (Myxobacterales). *Liebigs Ann. Chem.* **1983**: 1081–1095.
- Kim, Y. S., W. C. Bae, and S. J. Baek. 2003. Bioactive substances from myxobacteria. *Kor. J. Microbiol. Biotechnol.* **31**: 1–12.
- Krug, D., G. Zurek, O. Revermann, M. Vos, G. J. Velicer, and R. Müller. 2008. Discovering the hidden secondary metabolome of *Myxococcus xanthus*: A study of intraspecific diversity. *Appl. Environ. Microbiol.* **74**: 3058–3068.
- Kunze, B., N. Bedorf, W. Kohl, G. Höfle, and H. Reichenbach. 1989. Myxochelin A, a new iron-chelating compound from *Angiococcus disciformis* (Myxobacterales). Production, isolation, physico-chemical and biological properties. *J. Antibiot.* **42**: 14–17.
- Meiser, P., H. B. Bode, and R. Müller. 2006. The unique DKxanthene secondary metabolite family from the myxobacterium *Myxococcus xanthus* is required for developmental sporulation. *Proc. Natl. Acad. Sci. USA* **103**: 19128–19133.
- Newman, D. J., G. M. Cragg, and K. M. Snader. 2000. The influence of natural products on drug discovery. *Nat. Prod. Rep.* **17**: 215–234.
- Park, S., B. Lee, J. Kim, C. Lee, E. Jang, and K. Cho. 2004. Isolation and characterization of bacteriolytic wild myxobacteria. *Kor. J. Microbiol. Biotechnol.* **32**: 218–223.
- Reichenbach, H. and G. Höfle. 1993. Production of bioactive secondary metabolites, pp. 347–397. In M. Dworkin and D. Kaiser (eds.). *Myxobacteria*. American Society for Microbiology, Washington, DC.
- Reichenbach, H. and G. Höfle. 1999. Myxobacteria as producers of secondary metabolites, pp. 149–179. In S. Grabley and R. Thiericke (eds.). *Drug Discovery from Nature*. Springer Verlag, Berlin.
- Reichenbach, H. and M. Dworkin. 1992. The myxobacteria, pp. 3416–3487. In A. Balows, H. G. Trüper, M. Dworkin, W. Harder, and K.-H. Schleifer (eds.). *The Prokaryotes*, 2nd Ed. Springer-Verlag, New York.
- Rochfort, S. 2005. Metabolomics reviewed: A new “Omics” platform technology for systems biology and implications for natural products research. *J. Nat. Prod.* **68**: 1813–1820.
- Silas, G. V. B., S. Mas, M. Åkesson, J. Smedsgaard, and J. Nielsen. 2005. Mass spectrometry in metabolome analysis. *Mass Spect. Rev.* **24**: 613–346.
- Want, J. E., B. F. Cravatt, and G. Siuzdak. 2005. The expanding role of mass spectrometry in metabolite profiling and characterization. *Chembiochem* **6**: 1941–1951.